

Mort subite

Dr Laurent Fauchier

Cardiologie, CHU Trousseau, TOURS.

Objectifs

- Reconnaître les causes de mort subite (MS)
- (Changements dans la prise en charge en aigu de la mort subite)
- Comprendre la prise en charge diagnostique, pronostique et thérapeutique des arythmies à risque létal
- Mise à jour et perspectives sur l'utilisation du DAI dans la MS.

ACC/AHA/ESC 2006 guidelines for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death

Clinical Cardiology: New Frontiers

Sudden Cardiac Death

Douglas P. Zipes, MD; Hein J.J. Wellens, MD

Sudden cardiac death describes the unexpected natural death from a cardiac cause within a short time period, generally ≤ 1 hour from the onset of symptoms, in a person

patients in the convalescent phase after a large myocardial infarction experience sudden cardiac death in the year thereafter, overall they account for a small number of the total

Circulation 1998;98:2334-51.

Mort subite d'origine cardiaque

- Def : décès < 1 heure, hors de l'Hôpital, chez un patient préalablement stable
- Incidence : 400 000 – 500 000 / an aux U.S.A. (0.2 % / an ; 60 % du total des DC)
- Seulement 2% - 15% arrivent à l'hôpital
- > 50 % décèdent avant la sortie
- Taux de récurrence élevé

Causes of Sudden Cardiac Death

Ischemic Heart Disease

- CAD with MI or angina
- coronary artery embolism
- nonathogenic coronary artery disease
- coronary artery spasm

Nonischemic heart disease

- CAD without MI or angina
- cardiomyopathy - obstructive, nonobstructive, nonischemic
- valvular heart disease
- congenital heart disease
- prolonged QT syndrome
- preexcitation syndrome
- complete heart block
- arrhythmogenic RV dysplasia


Nonischemic Heart Disease (con't)

- myocarditis
- acute pericardial tamponade
- acute myocardial rupture

Noncardiac Disease

- sudden infant death syndrome
- drowning
- Pickwickian syndrome
- pulmonary embolism
- drug-induced
- airway obstruction
- no structural heart disease - primary electrical disease, chest wall trauma (commotio cordis), "Brugada's syndrome" (right bundle branch block and ST segment elevation V1 to V3)


Underlying Arrhythmia of Sudden Death




Adapted from Bayés de Luna A. Am Heart J. 1989;117:151-159.

Mechanisms of sudden death in heart failure

Ventricular tachycardia / Fibrillation (60 %)



Bradycardia - Electromechanical dissociation (40 %)




Intérêt du DAI


Traitement symptomatique
des troubles du rythme ventriculaire : TV et FV



VF Detection



continuous



VF Therapy
31 J. Shock



Sous -groupes à haut risque nécessitant évaluation

- Rescapés de mort subite
- Post-IDM, FE abaissée, et/ou ESV
- Syncopes répétées inexpliquées
- CMD idiopathique avec syncope ou TV
- CMH avec syncope ou TV
- Cardiomyopathie arythmogène VD
- Syndrome QT long

Evaluation of the survivor of sudden cardiac death

- Discontinue drugs (especially antiarrhythmics) if possible
- Correct metabolic and electrolyte abnormalities
- Evaluate for precipitating factors
- Optimize left ventricular function
- Control symptoms of active ischemia
- Systematic evaluation of antiarrhythmic drugs (Holter, EPS)
- Consider alternative therapy for drug-refractory patients

Methods of Evaluating Patients for Risk of Ventricular Arrhythmias


- History and physical
- 12-Lead ECG
- Signal-averaged ECG
- Holter monitor
- Echocardiogram
- Exercise test
- Cardiac catheterization
- Electrophysiology study

Le triangle de l'arythmogénèse

1. Substrat

2. Trigger

3. Modulateurs



Holter

Avantages

- largement accessible et non invasif
- interprétation facile
- facile à répéter au cours du suivi

Inconvénients

- valeur prédictive négative limitée
- critère d'efficacité d'une drogue ?
- différence entre l'effet d'une drogue (suppression ESV) et son efficacité (prévention de TVS ou FV)

Signal Averaged - ECG

ACC Expert Consensus

- Stratification du risque de survenue de TV soutenue en post-IDM avec QRS < 120 ms
- Identification des patients avec CMI et syncope inexpliquée qui ont une probabilité élevée de TV inductible.

Explo. Electrophysiologique

Avantages

- variabilité arythmie modeste
- peut documenter arythmie clinique et risque de MS
- localisation foyer arythmique

Inconvénients

- équipement, coût
- bcp de patients non inductible
- reproductibilité incertaine
- valeur de l'efficacité AA incertaine

Patient avec probabilité pre-test élevée d'EP + :

- ↓ **Fraction ejection VG**
- **TVNS**
- **PVT +**

Cardiopathies : marqueurs de MS

- CMI chronique ESV, TVNS, FEVG < 30%
- CMH Syncope, TVNS, hypoTA d'effort, ATCD fam de MS, HVG > 25-30 mm
- CMD TVNS (?), FEVG basse, VRS basse, QRS > 120 ms
- CMA VD Dilat. VD, TVNS, forme familiale
- HTA HVG, FEVG basse, Tb cond IV
- Amylose FEVG abaissée, Tb cond IV,
Sarcoidose ESV
Dystrophie & Myop

Cœur « sain » : Marqueurs de MS

Anomalies de repolarisation

- QT long QTc > 460 ms
- Familial Syncope, ATCD fam MS
- Acquis TV polym, TdP
- Brugada Syncope, ATCD fam MS, EP?


Préexcitation V

- WPW WPW, syncope, FA rapide


Clinical Substrates Associated with VF Arrest

- **Coronary artery disease**
- **Idiopathic cardiomyopathy**
- **Hypertrophic cardiomyopathy**
- **Long QT syndrome**
- **RV dysplasia**
- **Rarely: WPW syndrome**

VALIANT. Solomon et al, *NEJM* 2005




Annual Mortality in Heart Failure




Do all pts with HF require ICD to prevent sudden death?

Hsia et al. *Curr Control Trials Cardiovasc Med* 2000

Survival After Acute MI




Risk of Sudden Death: Data from GISSI-2 Trial



**Patients without
LV Dysfunction**

— No PVBs
- - 1-10 PVBs/h
..... > 10 PVBs/h



**Patients with
LV Dysfunction**

Prévention primaire de la MS



- **Beta bloqueurs**
- **IEC**
- **Revascularisation**
- **DAI**

Antiarrhythmic agents after MI CAST and SWORD Trials

Death or RCA

Death


Death




N Engl J Med 1991; 324:781

Lancet 1996; 348:7.


Outcome of EP-Guided Antiarrhythmic Drug Therapy in Sudden Death Survivors




MUSTT, NEJM 1999



ATMA, Lancet 1997



ATMA, Lancet 1997



Amiodarone


■ Toxicity

- Pulmonary fibrosis
- Hypo- or hyper-thyroidism
- Liver failure
- Bone marrow suppression
- Renal failure
- Photosensitivity
- Corneal deposits

■ Side effects

- Myalgias
- Gait disturbance
- Insomnia
- Prolongation of coagulation time (PT)
(need to reduce coumadin dosage)
- Digoxin toxicity (need to reduce digoxin dosage)

SCD Heft, *NEJM* 2005




No. at Risk

Amiodarone	845	772	715	484	280	97
Placebo	847	797	724	505	304	89
ICD therapy	829	778	733	501	304	103

CHF - STAT, *NEJM* 1995

Amiodarone in HF Patients with frequent ventricular arrhythmia.



OPTIC, JAMA 2006

	Beta-blocker (n = 138)	Amiodarone + Beta-blocker (n = 140)	Sotalol (n = 134)
<i>Any shock:</i>			
Rate/year (%)	38.5	10.3	24.3
Hazard ratio		0.27	0.61
95% CI		0.14-0.52	0.37-1.01
P value		< .001	.055

	Beta-blocker (n = 138)	Amiodarone + Beta-blocker (n = 140)	Sotalol (n = 134)
<i>Appropriate shock (%)</i>			
Rate/year (%)	22.0	6.7	15.1
Hazard ratio		0.30	0.65
95% CI		0.14-0.68	0.36-1.24
P value		.0035	.18
<i>Inappropriate shock</i>			
Rate/year (%)	15.4	3.3	9.4
Hazard ratio		0.22	0.61
95% CI		0.07-0.64	0.29-1.20
P value		.0055	.20

Ttt médical préventif de la mort subite

- Amiodarone : seul AA diminuant la mort subite (totale ??). Peut-être équivalent au DAI chez patient avec FE >35 %.
- Tous les autres AA sont inefficaces et/ou délétères pour diminuer la mortalité
- Traitements non AA apportant le plus de bénéfice pour réduire la mortalité :
 - Béta bloqueurs
 - IEC
 - Spironolactone
 - Huile de poisson (?)
 - Statines

Beta Blockers and Risk of Death in HF

Table 5. Summary of Mortality Across Trials and Best Comparison Subgroup*

	CIBIS-II	COPERNICUS	MERIT-HF	BEST	BCG
n	2647	2289	3991	2708	1668
Number of deaths	384 (14.5%)	320 (14.0%)	362 (9.1%)	860 (31.8%)	494 (30.0%)
Average length of follow-up (years)	1.3	0.87	1.0	2.0	2.1
Annual placebo mortality (%)	13.2	19.7	11.0	16.7	16.1
Hazard ratio and 95% confidence interval					
Mortality	0.66 (0.54–0.81) <i>P</i> < .0001	0.65 (0.52–0.81) <i>P</i> = .0014	0.66 (0.53–0.81) <i>P</i> < .0001	0.90 (0.78–1.02) <i>P</i> = .1024	0.77 (0.65–0.92) <i>P</i> = .0041
Cardiovascular death	0.71 (0.56–0.90) <i>P</i> = .0049	N/A	0.62 (0.50–0.78) <i>P</i> = .0003	0.86 (0.74–0.99) <i>P</i> = .041	0.71 (0.58–0.86) <i>P</i> = .0005
Sudden death	0.56 (0.39–0.80) <i>P</i> = .0011	N/A	0.59 (0.45–0.78) <i>P</i> = .0002	0.88 (0.72–1.07) <i>P</i> = .206	0.77 (0.59–0.999) <i>P</i> = .0488
Pump failure death	0.74 (0.48–1.14) <i>P</i> = .17	N/A	0.51 (0.33–0.79) <i>P</i> = .0023	0.85 (0.67–1.08) <i>P</i> = .188	0.64 (0.45–0.91) <i>P</i> = .0127

BEST, β -Blocker Evaluation of Survival Trial; BCG, BEST Comparison Subgroup; CIBIS-II, Cardiac Insufficiency Bisoprolol Study-II; COPERNICUS, Carvedilol Prospective Randomized Cumulative Survival Study; MERIT-HF, Metoprolol Intervention Trial in Congestive Heart Failure Trial; N/A, not applicable.

Domanski et al, *J Card Fail* 2003

Beta Blockers in SCD


The mode of action of beta blockers is not completely understood, but includes:

- Direct sympathetic blockade
- A membrane stabilizing effect
- A reduction in ischemic potential
- An increase in heart rate variability and baroreflex sensitivity
- A shorter corrected QT interval
- Prevention of stress-induced hypokalemia

IEC et prévention de MS en post IDM


Domanski et al, *JACC* 1999

	Sample Size		Sudden Cardiac Death		
	ACE	Placebo	ACE	Placebo	OR (95% CI)
Mortarino ²⁶	10	11	0	0	1.10 (0.02-60.30)
Oldroyd ²⁴	49	50	4	4	1.02 (0.26-4.02)
Nabel ¹⁷	20	18	0	1	0.29 (0.27-7.44)
Sharpe ²⁵	50	50	3	2	1.43 (0.27-7.61)
SMILE ¹⁴	772	784	4	11	0.39 (0.13-1.18)
EDP ²⁰	47	42	0	0	0.90 (0.02-46.09)
ECCE ¹⁸	104	104	1	0	3.03 (0.12-75.22)
CONSENSUS 2 ¹⁵	3044	3046	86	88	0.98 (0.72-1.32)
SAVE ⁵	1115	1116	105	125	0.83 (0.63-1.08)
AIRE ^{7,16}	1004	982	89	121	0.69 (0.52-0.93)
PRACTICAL ¹⁹	150	75	5	4	0.60 (0.17-2.16)
Søgaard ²¹	29	29	1	0	3.11 (0.12-79.43)
CATS ²²	149	149	3	4	0.77 (0.19-3.18)
TRACE ⁶	876	873	105	133	0.76 (0.58-1.00)
EDEN ²³	239	117	1	0	1.48 (0.06-36.56)
Overall	7658	7446	407	493	0.80 (0.70-0.92)



ICD Primary Prevention Trials

Extent of β -Blocker / ACE-Inhibitor Use
in Control Patients



GISSI-Prevenzione trial. *Lancet* 1999

	All (n=11 324)	Two-way analysis		
		n-3 PUFA (n=5666)	Control (n=5668)	Relative risk (95% CI)
Main endpoints				
Death, non-fatal MI, and non-fatal stroke	1500 (13.3%)	715 (12.6%)	785 (13.9%)	0.90 (0.82–0.99)
Cardiovascular death, non-fatal MI, and non-fatal stroke	1155 (10.2%)	547 (9.7%)	608 (10.8%)	0.89 (0.80–1.01)
Secondary analyses				
All fatal events	1017 (9.0%)	472 (8.3%)	545 (9.6%)	0.86 (0.76–0.97)
Cardiovascular deaths	639 (5.6%)	291 (5.1%)	348 (6.2%)	0.83 (0.71–0.97)
Cardiac death	520 (4.6%)	228 (4.0%)	292 (5.2%)	0.78 (0.65–0.92)
Coronary death	479 (4.2%)	214 (3.8%)	265 (4.7%)	0.80 (0.67–0.96)
Sudden death	286 (2.5%)	122 (2.2%)	164 (2.9%)	0.74 (0.58–0.93)
Other deaths	378 (3.3%)	181 (3.2%)	197 (3.5%)	0.91 (0.74–1.11)
Non-fatal cardiovascular events	578 (5.1%)	287 (5.1%)	291 (5.1%)	0.98 (0.83–1.15)

GISSI-Prevenzione trial. *Lancet* 1999

Pharmacological therapy	
Antiplatelet drugs	
Baseline	2601 (92.2%)
6 months	2308 (88.2%)
42 months	1707 (83.4%)
Angiotensin-converting-enzyme inhibitors	
Baseline	1298 (46.0%)
6 months	1033 (39.5%)
42 months	788 (38.5%)
β -blockers	
Baseline	1237 (43.9%)
6 months	1092 (41.7%)
42 months	807 (39.4%)
Cholesterol-lowering drugs	
Baseline	124 (4.4%)
6 months	782 (28.6%)
42 months	1003 (46.0%)
<hr/>	
Revascularisation procedures*	
CABG or PTCA	
Baseline	135 (4.8%)
6 months	433 (15.3%)
42 months	689 (24.3%)

Revascularisation


- La revascularisation (par pontage) ne diminue pas l'incidence de récurrence de TV / FV sauf en cas de relation directe documentée d'un événement avec l'ischémie.

CASS Registry Circulation 1986


1
trc

2
trc


3
trc



CASS : Coronary Artery Surgery Study



Secondary Prevention Trials: Reduction in Overall Mortality with ICD Therapy




¹ The AVID Investigators. *N Engl J Med.* 1997;337:1576-83.

² Kuck K. *Circ.* 2000;102:748-54.

³ Connolly S. *Circ.* 2000;101:1297-1302.

Secondary Prevention Trials: Reduction in Mortality with ICD Therapy



¹ The AVID Investigators. *N Engl J Med.* 1997;337:1576-83.


² Kuck K. *Circ.* 2000;102:748-54.

³ Connolly S. *Circ.* 2000;101:1297-1302.

“Climat” au début des études de prévention primaire avec DAI (1990...)

- Incertitude sur le bénéfice du DAI (vs chirurgie des TV, EEP sériées sous AA...)
- DAI épiscardique (mortalité op. 4-5 %)
- Etudes de prévention secondaire non publiées
- Etudes de prévention primaire jugées non éthiques par beaucoup


Sudden Cardiac Death Incidence and Total Events



MADIT / MUSTT / MADIT II / CABG-Patch


Patient Inclusion Criteria	MADIT	MUSTT	MADIT-II	CABG-Patch
CAD/Post-MI	X	X	X	X
Low LVEF	X ($\leq 35\%$)	X ($\leq 40\%$)	X ($\leq 30\%$)	X ($\leq 36\%$)
SAECG				X
NSVT	X	X		
Inducible VT on EPS	X	X		
Inducible, non-suppressible VT on EPS	X			

CABG Patch Trial (*Bigger Jr, NEJM 1997*)




Defibrillator group	446	384	313	213	61
Control group	454	389	308	199	57

CABG Patch Trial (*Bigger Jr, NEJM 1997*)




ICD Benefit is ~ Double In Primary Prevention Compared to Secondary



SCD HeFT: Design

- Enrollment Criteria:
 - Ejection Fraction <36%
 - Ischemic or idiopathic Dilated Cardiomyopathy
 - Class II, III CHF
- Randomization:
 - Standard CHF Therapy + Placebo
 - Standard CHF Therapy + Amiodarone
 - Standard CHF Therapy + ICD
- Statistical Assumptions:
 - Mortality with conventional therapy 25%
 - Planned enrollment 2500 pts; fixed sample size
 - 2.5 yr follow-up

SCD Heft, *NEJM* 2005



No. at Risk

Amiodarone	845	772	715	484	280	97
Placebo	847	797	724	505	304	89
ICD therapy	829	778	733	501	304	103

Sanders,
NEJM
2005

Table 1. Health and Economic Outcomes of the Prophylactic Implantation of an ICD as Compared with Control Therapy.*

Trial and Strategy	Cost	Increase in Cost Related to ICD	Life Expectancy	Increase in Life Expectancy Related to ICD	QALY	Increase in QALY Related to ICD	Incremental Cost-Effectiveness			
							\$/Life-Yr	Baseline Efficacy	High Efficacy \$/QALY	
	\$			year						
MADIT I										
Control	38,300		4.06		2.98					
ICD	130,400	92,100	7.70	3.64	5.62	2.64	25,300	34,900	27,000	
CABG Patch										
Control	78,600		8.41		6.13					
ICD	134,400	55,700	8.01	(0.40)	5.84	(0.29)	Dominated	Dominated	84,200	
MUSTT										
Control	44,300		4.72		3.46					
ICD	145,800	101,500	8.86	4.14	6.45	2.99	24,500	34,000	28,800	
MADIT II										
Control	57,500		6.16		4.51					
ICD	136,900	79,400	8.20	2.03	5.98	1.47	39,000	54,100	37,200	
DEFINITE										
Control	84,400		9.03		6.57					
ICD	184,900	100,500	11.75	2.73	8.53	1.96	36,800	51,300	34,500	
DINAMIT										
Control	88,300		9.44		6.87					
ICD	147,200	58,800	8.96	(0.48)	6.53	(0.34)	Dominated	Dominated	70,900	
COMPANION										
Control	37,800		4.01		2.95					
ICD	106,100	68,300	5.88	1.87	4.31	1.36	36,500	50,300	36,100	
SCD-HeFT										
Control	57,800		6.19		4.53					
ICD	128,800	71,000	7.59	1.40	5.54	1.01	50,700	70,200	45,600	

Données épidémiologiques

Insuffisance cardiaque

Insuffisance Cardiaque

Prévalence 1 à 2 %



Mortalité annuelle

12 - 15 % classe I - 60 % classe IV




Mortalité rythmique

50 à 60 % classe I - II - 20 à 30 % classe IV

SCD Heft, *NEJM* 2005


A NYHA Class II



No. at Risk

Amiodarone	601	563	536	378	222	76
Placebo	594	563	522	367	218	72
ICD therapy	566	550	531	371	236	80

B NYHA Class III




No. at Risk

Amiodarone	244	209	179	106	58	21
Placebo	253	234	202	138	86	17
ICD therapy	263	228	202	130	68	23

SCD Heft, *NEJM* 2005

A Ischemic CHF


	Hazard Ratio (97.5% CI)	P Value
Amiodarone vs. placebo	1.05 (0.81–1.36)	0.66
ICD therapy vs. placebo	0.79 (0.60–1.04)	0.05



No. at Risk						
Amiodarone	426	384	346	227	130	46
Placebo	453	415	370	244	152	48
ICD therapy	431	395	365	244	144	48

B Nonischemic CHF

	Hazard Ratio (97.5% CI)	P Value
Amiodarone vs. placebo	1.07 (0.76–1.51)	0.65
ICD therapy vs. placebo	0.73 (0.50–1.07)	0.06



No. at Risk						
Amiodarone	419	388	369	257	150	51
Placebo	394	382	354	261	152	41
ICD therapy	398	383	368	257	160	55

Other Noninvasive Evaluations

- **QT or JT dispersion** - measured as interlead variability of QT or JT intervals on the surface ECG
- **Repolarization (T wave) alternans (RPA)** - refers to a variability in the timing or morphology of repolarization that occurs in alternate beats on the surface electrocardiogram
- **Heart rate (RR) variability** - measurements provide a low-cost, widely-available method for assessing the status of the parasympathetic/sympathetic nervous system

Identification des patients à risque de mort subite dans l'insuffisance cardiaque

Paramètres	Analyse univariée	Multivariée	Controversée
Maladie coro.	X	X	X
Syncope	X	X	
TVS/FV doc.	X		
TVNS	X	X	X
Variabilité sinusale	X	X	X
Dispersion QT	X	X	X
Dispersion QRS	X	X	
Alternance onde T	X	X	
Hyponatrémie	X	X	X
FeVG	X	X	X
Index cardiothorac.	X		

DINAMIT

- . IDM < 6-30 j, FE VG < 35%,
- . FC > 80/ ' ou SDNN < 70ms,

ttt med_


(IEC, Beta B...)

ttt med

+ Def AI

- . Critère principal : mortalité globale
- Critères 2^{aires} : coûts, qualité de vie,...
- . N > 625, suivi > 2 ans


DINAMIT - NEJM 2004



No. at Risk

ICD group	315	299	258	211	172	123	82	25
Control group	318	305	272	217	172	124	79	31

COMPANION, 2003



Implantable Cardioverter Defibrillator in High-Risk Long QT Syndrome Patients

Zareba et al JCE 2003; 14:337


- **125 LQTS patients with ICDs: 54 with CA and 19 with recurrent syncope despite BB = 73**
- **Compared to 161 LQTS patients with similar indications but no ICD: 89 CA, 72 syncope**
- **1 death in ICD group (1.3%) over 3 years, and 26 deaths (16%) in non-ICD group over 8 years (p=.07)**

Defibrillator v. β -Blockers for Unexplained Death in Thailand (DEBUT)

Nademanee et al *Circulation* 2003; 107:2221


- 86 survivors of SUDS
- Randomized to ICD v beta blocker
- 3 year follow up had 7 deaths in beta blocker group, none in ICD (p=.02)
- 12 ICD patients received shocks for VF

Overall U.S. ICD Utilization



Ruskin JN, Hallstrom AP, Zipes DP, Camm AJ, McGrory-Ussat ME. ICD Utilization Based on Discharge Diagnoses from Medicare and Managed Care Patients J Cardiovasc Electrophysiol 13:38-43, 2002

Estimated ICD Utilization



Conclusions

- Le DAI est plus efficace que le traitement médical pour diminuer le risque de mort subite
- Le bénéfice est particulièrement net en cas de FEVG basse
- Le DAI reste sous-utilisé
- Baisse des prix du DAI pour élargir son utilisation ?
- Définir des populations ciblées qui bénéficient du DAI ?



- M 68, aéroport Indianapolis
- FdRCV : Tabac, HTA




- Appel 911 : 0' 54"



- Manœuvres Ressuscitation : 2' 10"



- Défibrillation : 3' 43"





- Réveil J1
- Coro : 3 tronculaire
- + dysfonction VG
- PAC + amiodarone
- Sortie J14
- DAI J45
- Reprise W 6h / j à J60

Mais ...

- **80 % à 90 % des arrêts cardiaques extra-hospitaliers surviennent à domicile**
- **De l'intérêt d'une prévention primaire performante.**